



PRODUCT SPECIFICATION

biolla DRIED GLUCOSE SYRUP DE 30

Dextrose Equivalent 28.0-35.0 / Grade: EP 11.2

Product description

Molecular Formula:	$H(C_6H_{10}O_5)_n$ -OH
EI N ECS:	232-940-4
CAS No:	9050-36-6
Appearance:	homogeneous free flowing fine powder, microgranules
Colour:	white
Taste and odour:	typical to maltodextrins, without foreign taste and odour
Intended use:	intended for use as a carbohydrate component, structure- forming agent, forming agent, natural sweetener, sweetener regulator, stabilizer, neutral carrier of flavouring additives in various industries.
Production method:	partial starch cleavage product consisting of a multicomponent mixture of glucose, maltose, maltotriose and polysaccharides in various ratios.

- Plant based product produced from corn grain not containing GMO.
- Complies to the Food Chemical Codex and to the European / US Pharmacopeia monographs on Maltodextrin.

Nutrition value in 100 g of a product, min.	93/100 g
Carbohydrates, %, min	93
Fats, %	0
Proteins, %	0
Energy value (calorific value) kcal / kJ per 100g maltodextrin, min	372/1581
Composition	Dried Glucose Syrup 100%

BIC/SWIFT: COBADEFFXXX

03 1796 06 biolla.de/agb EHHXXX







Parameter	Units	Typical value	Method of analysis
Moisture, max	%	5.0	Determination of loss in mass on drying (130°C / 5 g / 1,5 h) (Ref. ISO 1666:1996)
Dextrose Equivalent (Lane-Eynon method), max	%	28.0 – 35.0	Titremetry, Lane-Eynon method (Ref. ISO 5377:1981)
Sulphated Ash Content, %, max	%	0.20	Weighing of residuals from burning a weighed portion of maltodextrin in muffle oven at 550°C $(+H_2SO_4)$
			(Ref. ISO5809:1981)
Bulk density (loose)	kg/dm ³	0.45 - 0.60	Internal method
			(Product weight in 1dm ³ in free state)
Sulphur dioxide content (SO ₂), max	mg/kg	10.0	lodometry (Oxidation of sulphurous acid in a 50% solution of maltodextrin in the presence of a starch indicator)
Solubility, min	%	98.0	Internal method (Dissolving 50 g of dried glucose syrup in 100 g of hot water)
pH value	pH units	4.5-6.0	pH-metry
			40 % solution of dried glucose syrup
Particles size:	%	5.0	Granulometry
Residue on sieve 200 µm, max			
Particles size:	%	90.0	Granulometry
Residue on sieve 40 µm, min			-

Carbohydrate composition (in DS)

Glucose (DP ₁)	%	3-7	HPLC (Ref. ISO 10504)
Maltose (DP ₂)	%	12-18	HPLC (Ref. ISO 10504)
Maltose (DP ₃)	%	20-25	HPLC (Ref. ISO 10504)
Higher sugars (DP ₄₊)	%	On balance	HPLC (Ref. ISO 10504)

BIC/SWIFT: COBADEFFXXX

biolla.de/agb





Food safety parameters

Total Aerobic and Anaerobic Microbial Count, max	CFU/g	50.0	Ref. ISO 4833
Yeast, max	CFU/g	50.0	Ref. ISO 7954
Molds, max	CFU/g	50.0	Ref. ISO 7954
Coliforms	CFU/1.0 g	not allowed	Ref. ISO 4831, ISO 4832
Pathogenic microbes including Salmonella	CFU/25 g	not allowed	Ref. EN ISO 6579-1

Content of Heavy Metal (mg/kg), max

Lead (Pb)	0.10		
Cadmium (Cd)	0.10	Atomic absorptive method	
Arsenic (As)	0.10	(mineralization and determination with atomic absorptive spectrometer)	
Mercury (Hg)	0.02	, and the part of	

Content of GMO (PCR Real - Time)

Genetically Modified Organisms	not allowed	PCR Real – Time
		(Ref. ISO 21569, ISO 21571)

Additional information

Shelf life is 2 years. Dried Glucose Syrup must be stored at relative humidity not exceeding 75% Standard packing – 25 kg paper bags with PE lining.

BIC/SWIFT: COBADEFFXXX

biolla.de/agb

